



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd



Document heading

doi:10.1016/S2222-1808(14)60327-3

© 2014 by the Asian Pacific Journal of Tropical Disease. All rights reserved.

Prevalence of fasciolosis and dicrocoeliosis in slaughtered sheep and goats in Amol Abattoir, Mazandaran, northern Iran

Ali Khanjari^{1*}, Alireza Bahonar¹, Sepideh Fallah¹, Mahboube Bagheri¹, Abbas Alizadeh², Marjan fallah³, Zahra Khanjari⁴

¹Department of Food Hygiene, Faculty of Veterinary Medicine, University of Tehran, P.O. Box 14155–6453, Tehran, Iran

²Department of Physiology, Faculty of Veterinary Medicine, Islamic Azad University, Science and Research Branch, Tehran, Iran

³Faculty of Veterinary Medicine, University of Urmia, Urmia, Iran

⁴Faculty of Medicine, Sari Branch, Islamic Azad University, Mazandaran, Iran

PEER REVIEW

Peer reviewer

Hossein Esmaeili, Assistant professor, Department of Microbiology, Faculty of Veterinary Medicine, University of Tehran, Iran.

Tel:00982161117051

Fax: 00982166933322

E-mail: hesmaeli@ut.ac.ir

Comments

This manuscript contains useful information with regard to an important parasitic disease which is responsible for considerable economic losses in developing countries. In my point of view, Authors have been successful to demonstrate the relationship between the prevalence of the liver fluke and the sex/ season/ specie factor.

Details on Page 123

ABSTRACT

Objective: The liver flukes, *Fasciola* spp. and *Dicrocoelium dendriticum*, infect ruminants and other mammalian extensively and cause major diseases of livestock that produce considerable economic losses.

Methods: A survey of 2391 sheep and goats slaughtered at an abattoir in Amol region, northern Iran was used to determine the prevalence of the liver flukes infection based on season, sex and specie of the animals.

Results: The results revealed that the prevalence rate of *Fasciola* spp. and *Dicrocoelium dendriticum* was 6.6% and 4.3% respectively. Dicrocoeliosis was more dominant in female animals (7.1%) whereas there was no sex-related difference in the prevalence of *Fasciola* spp. in male and female animals. Furthermore, Fasciolosis was significantly more prevalent than dicrocoeliosis in both sheep and goats. The Seasonal prevalence of *Fasciola* spp. was highest ($P < 0.005$) during spring (8.3%) followed in order by autumn (8.1%), winter (5.9%) and summer (4.0%) but Dicrocoeliosis did not follow any seasonal pattern.

Conclusions: According to this study, it can be concluded that Amol is regarded as an endemic region for *Fasciola* spp and *D. dendriticum* infection. Moreover, *Fasciola* spp. is the most widespread liver fluke found in sheep and goats which is more dominant in sheep than goats.

KEYWORDS

Liver fluke, Dicrocoeliosis, Fasciolosis, Sheep, Goat, Abattoir

1. Introduction

The liver flukes are recognized as one of the most important ruminants helminthic parasites which are found in many parts of the world [1,2]. *Fasciola* spp. and

Dicrocoelium dendriticum (*D. dendriticum*) are the common liver flukes in Iran. The principal definitive hosts of these parasites are cattle, sheep and goat. However, certain other mammals, including humans, may be infected as an accidental host [3,4]. Due to the increasing number of

*Corresponding author: Ali Khanjari, Department of Food Hygiene, Faculty of Veterinary Medicine, Tehran University, P.O.Box: 14155–6453, Tehran, Iran.

Tel: +98–21–61117023

Fax: +98–21–66933222

E-mail: khanjari@ut.ac.ir

Fundation Project: Supported by the research council of University of Tehran (Grant No. 28747/1/6).

Article history:

Received 1 Dec 2013

Received in revised form 4 Dec, 2nd revised form 10 Dec, 3rd revised form 16 Dec 2013

Accepted 15 Jan 2014

Available online 28 Apr 2014

human cases, the liver flukes should be considered as an emerging public health concern[5]. On the other hand, they cause major diseases of livestock that produce important economic losses due to mortality, liver condemnation, reduced production of meat, milk, and wool, and expenditures of anthelmintics[6].

Fasciola spp. has an indirect life cycle involving domestic and wild herbivorous mammals and humans as definitive hosts, and freshwater gastropods of the family Lymnaeidae as intermediate hosts[7]. *Fasciola hepatica* (*F. hepatica*) occurs in temperate areas, while *Fasciola gigantica* mainly occurs in tropical zones, but both species overlap in subtropical areas[8]. Temperature, rainfall and soil moisture influence the activity and abundance of the intermediate hosts of *F. hepatica* and *Fasciola gigantica*. Consequently, current climatic conditions can be influential in the development of Fasciolosis[9].

In Iran, fasciolosis is present in many provinces including Kurdistan, Zanjan, Kermanshah, Mazandaran, Tehran, Azerbaijan, Gilan, Fars and Khuzestan[10]. Although the prevalence of fasciolosis among domestic animals is higher in the southern part of the country, the number of reported human disease cases is significantly higher in the Northern provinces situated along the shore of the Caspian Sea[11]. Both *Fasciola hepatica* and *Fasciola gigantica* are endemic in the Northern regions of Iran[10,11,12].

Dicrocoeliasis, a disease of grazing ruminants, is less severe than fasciolosis. While the economic losses, mainly as a result of affected liver condemnation, are considerable[13]. Two intermediate hosts are necessary to complete parasite development: a land snail (*Zebrina* sp., *Helicella* sp., *Cionella* sp.) and an ant (*Formica* sp., *Lasius* sp.). The Final host infection occurs by ingesting the infected ants[14]. Unlike *Fasciola* spp., the intermediate hosts of *D. dendriticum* do not require a moist environment and are widely present in pastures [15].

This study was conducted to estimate the prevalence of the liver flukes infection in sheep and goats slaughtered in Amol abattoir, Mazandaran, Iran, based on season, sex and specie of the animals.

2. Materials and methods

This survey was conducted from September 23, 2010 to September 23, 2011 at an abattoir in Amol region, Mazandaran province, Iran. Amol is located in the middle of the Mazandaran province which is in the north of Iran and has a borderline humid subtropical/Mediterranean climate.

A total number of 2391 sheep and goats, presented to be slaughtered at the abattoir, were randomly sampled during 4 seasons. Specie and sex information were documented

before inspection, on the basis of physical appearance of each sample, then the livers were inspected according to the method described by Ogambo–Ongoma (1972) to recognize fasciolosis and dicrocoeliasis[16]. The parasites were identified by their morphological characteristics.

The recorded data, acquired by visualization, palpation and incision of livers, was used to extract the prevalence rate of these parasites. The prevalence rate was sorted seasonally to determine the association of infection rate and seasons. Analysis of data was done, using SPSS software (Version 6.0). Seasonal pattern was investigated with Chi-square (χ^2) test and P-value less than 0.05 considered statistically significant.

3. Results

According to the results, among 2391 sheep and goats sampled during the study, 157 (6.6%) and 104 (4.3%) were infected by *Fasciola* spp. and *D. dendriticum* respectively. Dicrocoeliasis was recognized in 7.1% of female animals, whereas only 2% of male animals were infected by *D. dendriticum*. The difference of *D. dendriticum* prevalence in relation to the animals sex was statistically significant ($P < 0.005$). However, there was no sex-related difference in the prevalence of *Fasciola* spp. in male and female animals ($P > 0.1$). More details are shown in Table 1.

Table 1
Sex-wise prevalence of *Fasciola* spp. and *Dicrocoelium dendriticum* in sheep and goats slaughtered in Amol region abattoir

Liver fluke	Male animals			Female animals			χ^2	P
	Slaughtered	Infected	Infected	Slaughtered	Infected	Infected		
	No.	No.	%	No.	No.	%		
<i>Fasciola</i> spp.	1287	13	1.0%	1104	144	13.0%	1.403	0.2362
<i>D. dendriticum</i>	1287	26	2.0%	1104	78	7.1%	36.354	0.000

The results revealed that 7.7% and 5.4% of the sheep and goats were infected by *Fasciola* spp. respectively. On the other hand, dicrocoeliasis was recognized in 5.7% of the sheep and 3.0% of the goats. Thus, fasciolosis was significantly more prevalent than dicrocoeliasis in both sheep and goats (Table 2).

Table 2
Prevalence of *Fasciola* spp. and *Dicrocoelium dendriticum* in sheep and goats slaughtered in Amol region abattoir

Liver fluke	Sheep			Goats			χ^2	P
	Slaughtered	Infected	Infected	Slaughtered	Infected	Infected		
	No.	No.	%	No.	No.	%		
<i>Fasciola</i> spp.	1215	93	7.7%	1176	64	5.4%	4.767	0.0290
<i>D. dendriticum</i>	1215	69	5.7%	1176	35	3.0%	10.493	0.0011

Furthermore, the prevalence of fasciolosis and dicrocoeliasis was evaluated during four seasons of the year and as it is shown in Table 3, the *Fasciola* spp. seasonal prevalence was highest during spring (8.3%) followed in

order by autumn (8.1%), winter (5.9%) and summer (4.0%) ($P=0.0071$). Dicrocoeliasis prevalence did not follow any seasonal pattern, and seasonal differences were not statistically significant ($P=0.6607$).

Table 3

seasonal prevalences of *Fasciola* spp. and *Dicrocoelium dendriticum* in sheep and goats slaughtered in Amol region abattoir

Season	Slaughtered No.	<i>Fasciola</i>	<i>Fasciola</i>	<i>D.</i>	<i>D.</i>
		spp. Infected No.	spp. Infected %	<i>dendriticum</i> Infected No.	<i>dendriticum</i> Infected %
Spring	601	50	8.3%	30	5.0%
Summer	599	24	4.0%	27	4.5%
Autumn	595	48	8.1%	21	3.5%
Winter	596	35	5.9%	26	4.4%
Total	2391	157	6.6%	104	4.3%

4. Discussion

This study was carried out during a one-year period at an abattoir in Amol region, Mazandaran province in the North of Iran. Previous studies in Asia have shown that the liver flukes occur in various countries including Iraq, Pakistan, Turkey, Saudi Arabia, Tunisia, Bangladesh, India, China, Japan and Korea[17,18,19,20,21,22,23,24,25,26,27,28]. Moreover, there are numerous reports of fasciolosis and dicrocoeliasis from different parts of Iran including Fars, Mazandaran, Azarbaijan, Tehran, Khorasan and Gilan provinces[11,12,29,30,31,32,33].

In the current study, the prevalence of *Fasciola* spp. in sheep and goats was 7.7% and 5.4% respectively. In a study in 2004 at Amol city abattoir, Fasciolosis rate was 5.8% for sheep and 0.5% for goats[12]. On the other hand, Oryan *et al.* in 2011 reported that The infection rate due to *Fasciola* spp. was 0.35% in the sheep and 0.2% in the goats in Northeastern Iran[32]. As with the development of the intermediate host, temperature (>9.5 °C), rainfall and soil moisture are important factors influencing the development of the parasite from egg to miracidium. Thus, regional and annual variations in the prevalence of *F. hepatica* are associated with prevailing weather conditions[9]. In regard to host species, Fasciolosis was significantly more prevalent in sheep than goats. Both sheep and goats are very susceptible to *F. hepatica*, have no resistance to reinfection[7]. Higher prevalence in sheep may be due to the grazing habits of the animal species; goats graze on leaves and branches on bushes and trees but sheep graze on plants on the ground where metacercaria are mostly found. So, the possibility of infection with metacercaria is higher in sheep than goats[34].

According to the results, the occurrence rate of *D. dendriticum* was 5.7% in sheep and 3.0% in goats. Dicrocoeliasis prevalence has been reported 4.53% and 1.4% in northeast of Iran[32]and 2.4% and 1.85% in Mazandaran province[35]in sheep and goats respectively. The infection

has been described in either lowlands or mountain pastures at high altitudes and it appears that chalk or alkaline soils favour the development of the snail intermediate hosts and ants[36]. In addition, Cringoli *et al.* showed that large pastures and the presence of streams/brooks in pasture may be risk factors for *D. dendriticum* infection [37]. Sheep seem to be more susceptible to *D. dendriticum* than goats[38], which is in agreement with the results of the present study. Our results could not support any association between fasciolosis rate and sex of the animals which is observed by other investigators[39]. However, *D. dendriticum* was significantly more prevalent in female animals. Females show a higher dicrocoeliasis prevalence than males, probably as a result of different grazing habits[38]. On the other hand, it is assumed that sex is a determinant influencing prevalence of parasitism and females are more prone to parasitism during pregnancy and peri-parturient period due to stress and decreased immune status[18].

The prevalence of *Fasciola* spp. showed significant seasonal differences in this study. The hatching of fluke eggs and the multiplication of the snail intermediate host require high rainfall and temperatures (>10 °C) [40]. These conditions generally occur in the spring and autumn, when many fluke eggs hatch, snails multiply and then cercariae develop and are released on wet pastures before encysting onto herbage. The seasonality pattern in fasciolosis prevalence has been also observed by other investigators[41,42].

There are two important features that differentiate the epidemiology of *D. dendriticum* from that of *Fasciola* spp. Firstly, unlike *Fasciola* spp., the intermediate hosts of *Dicrocoelium* do not require a moist environment and are widely present in pastures and secondly, the fluke eggs can survive for months on these pastures[40]. Seasonality of this infection is favored by movement of the animals from lowland to mountain pastures where they become infected by the ants and then bring the infection back to the valley during the winter [38]. However, no significant correlation between *D. dendriticum* prevalence and seasons was detected in this study.

This abattoir survey generally reflected the disease situation in the Amol region. According to this study, it can be concluded that Amol is regarded as an endemic region for *Fasciola* spp and *D. dendriticum* infection. Moreover, *Fasciola* spp. is the most widespread liver fluke found in sheep and goats which is more dominant in sheep than goats. More surveys are suggested to be carried out to collect more data about the liver flukes infection prevalence and risk factors for developing a prediction model in small ruminants in northern Iran. The identified risk factors and the prediction model can be useful to formulate appropriate control strategies and decrease the economic loss due to

condemnation of infected livers.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

The authors would like to thank the authorities of faculty of veterinary medicine, University of Tehran for their financial support and cooperation. This Survey was financially supported by the research council of University of Tehran (grant number: 28747/1/6).

Comments

Background

Fasciolosis and dicrocoeliosis are common parasitic diseases of ruminants in Iran, making significant economic losses due to weight loss, reduction of milk production and condemned livers. Both parasites have indirect life cycle, thus various factors such as pasture habits, climate conditions may affect the prevalence.

Research frontiers

This study reports the prevalence of common liver flukes (dicrocoeliosis and fasciolosis) in a northern province of Iran located next to the Caspian sea in relation to season, sex and species of 2391 sheep and goats presented to the slaughter throughout one year.

Related reports

In the regions with similar climate condition to Iran, Fasciolosis is more prevalent than dicrocoeliosis. According to the intermediate host which needs moist to complete the life cycle, Fasciolosis prevalence shows more seasonal pattern than dicrocoeliosis. Sex related differences are less reported.

Innovations and breakthroughs

Collecting data about the liver fluke occurrence in sheep and goats in a region simultaneously may help to highlight the specie regarded differences. Also little data is reported about liver fluke prevalence in goats and seasonal pattern differences in Iran.

Applications

New prevalence data from the different regions may help

to understand the factors affecting the occurrence of the disease and improve control strategies.

Peer review

This manuscript contains useful information with regard to an important parasitic disease which is responsible for considerable economic losses in developing countries. In my point of view, Authors have been successful to demonstrate the relationship between the prevalence of the liver fluke and the sex/ season/ specie factor.

References

- [1] Kantzoura V, Kouam MK, Feidas H, Teofanova D, Theodoropoulos G. Geographic distribution modelling for ruminant liver flukes (*Fasciola hepatica*) in south–eastern Europe. *Int J Parasitol* 2011; **41**(7): 747–753.
- [2] Massoud AM, Shalaby HA, El Khateeb RM, Mahmoud MS, Kutkat MA. Effects of Mirazid and myrrh volatile oil on adult *Fasciola gigantica* under laboratory conditions. *Asian Pac J Trop Biomed* 2012; **2**(11): 875–884.
- [3] Conceição M, Durão R, Costa I, Castro A, Louza A, Costa J. Herd–level seroprevalence of fasciolosis in cattle in north central Portugal. *Vet Parasitol* 2004; **123**(1): 93–103.
- [4] McCann CM, Baylis M, Williams DJ. The development of linear regression models using environmental variables to explain the spatial distribution of *Fasciola hepatica* infection in dairy herds in England and Wales. *Int J Parasitol* 2010; **40**(9): 1021–1028.
- [5] Nguyen TG, Van De N, Vercruyse J, Dorny P, Le TH. Genotypic characterization and species identification of *Fasciola* spp. with implications regarding the isolates infecting goats in Vietnam. *Exp parasitol* 2009; **123**(4): 354–361.
- [6] Kleiman F, Pietrokovsky S, Prepelitchi L, Carbajo AE, Wisnivesky–Colli C. Dynamics of *Fasciola hepatica* transmission in the Andean Patagonian valleys, Argentina. *Vet Parasitol* 2007; **145**(3): 274–286.
- [7] Issia L, Pietrokovsky S, Sousa–Figueiredo J, Stothard JR, Wisnivesky–Colli C. *Fasciola hepatica* infections in livestock flock, guanacos and coypus in two wildlife reserves in Argentina. *Vet Parasitol* 2009; **165**(3–4): 341–344.
- [8] Alasaad S, Soriguer RC, Abu–Madi M, El Behairy A, Jowers MJ, Baños PD, et al. A TaqMan real–time PCR–based assay for the identification of *Fasciola* spp. *Vet Parasitol* 2011; **179**(1–3): 266–271.
- [9] Relf V, Good B, Hanrahan J, McCarthy E, Forbes A, deWaal T. Temporal studies on *Fasciola hepatica* in *Galba truncatula* in the west of Ireland. *Vet Parasitol* 2011; **175**(3–4): 287–292.
- [10] Amor N, Halajian A, Farjallah S, Merella P, Said K, Slimane BB. Molecular characterization of *Fasciola* spp. from the endemic area of northern Iran based on nuclear ribosomal DNA

- sequences. *Exp Parasitol* 2011; **128**(3): 196–204.
- [11] Ashrafi K, Valero MA, Panova M, Periago MV, Massoud J, Mas-Coma S. Phenotypic analysis of adults of *Fasciola hepatica*, *Fasciola gigantica* and intermediate forms from the endemic region of Gilan, Iran. *Parasitol Int* 2006; **55**(4): 249–260.
- [12] Moghaddam A, Massoud J, Mahmoodi M, Mahvi A, Periago M, Artigas P, et al. Human and animal fascioliasis in Mazandaran province, northern Iran. *Parasitol Res* 2004; **94**(1): 61–69.
- [13] Ahmadi R, Sikejor EM, Maleki M. Prevalence of *D. dendriticum* infection in cattle, sheep and goat in Gilan province, Northern Iran. *J Anim Vet Adv* 2010; **9**(21): 2723–2724.
- [14] Le Bailly M, Bouchet F. Ancient dicrocoeliosis: Occurrence, distribution and migration. *Acta Trop* 2010; **115**(3): 175–180.
- [15] Otranto D, Traversa D. A review of dicrocoeliosis of ruminants including recent advances in the diagnosis and treatment. *Vet Parasitol* 2002; **107**(4): 317–335.
- [16] Ogambo–Ongoma AH. Fascioliasis survey in Uganda. *Bull Epizoot Dis Afr* 1972; **20**(1): 35–41.
- [17] Ridha AM, Al–Sadi HI. Pathology of the digestive system of goats in northern Iraq. *Small Ruminant Res* 1992; **9**(1): 79–91.
- [18] Khan MN, Sajid MS, Khan MK, Iqbal Z, Hussain A. Gastrointestinal helminthiasis: prevalence and associated determinants in domestic ruminants of district Toba Tek Singh, Punjab, Pakistan. *Parasitol Res* 2010; **107**(4): 787–794.
- [19] Farooq Z, Mushtaq S, Iqbal Z, Akhtar S. Parasitic helminths of domesticated and wild ruminants in Cholistan desert of Pakistan. *Int J Agric Biol* 2012; **14**(1): 63–68.
- [20] Kara M, Gicik Y, Sari B, Bulut H, Arslan MO. A slaughterhouse study on prevalence of some helminths of cattle and sheep in Malatya province, Turkey. *J Anim Vet Adv* 2009; **8**(11): 2200–2205.
- [21] Nasher AK. Parasites of livestock in Asir province, southwestern Saudi Arabia. *Vet Parasitol* 1990; **37**(3–4): 297–300.
- [22] Hammami H, Hamed N, Ayadi A. Epidemiological studies on *Fasciola hepatica* in Gafsa oases (south west of Tunisia). *Parasite* 2007; **14**(3): 261–264.
- [23] Hossain MM, Paul S, Rahman MM, Hossain FMA, Hossain MT, Islam MR. Prevalence and economic significance of caprine fascioliasis at Sylhet district of Bangladesh. *Pak Vet J* 2011; **31**(2): 113–116.
- [24] Kumari S, Sinha SR, Sinha S, Hoda MZ, Mandal KG, Sharma SK. Incidence of gastrointestinal helminthosis in sheep and goats in patna (Bihar). *J Vet Parasitol* 2010; **24**(1): 97–99.
- [25] Lone BA, Chishti MZ, Ahmad F, Tak H. A survey of gastrointestinal helminth parasites of slaughtered sheep and goats in Ganderbal, Kashmir. *Global Vet* 2012; **8**(4): 338–341.
- [26] Huang W, He B, Wang C, Zhu X. Characterisation of *Fasciola* species from Mainland China by ITS–2 ribosomal DNA sequence. *Vet Parasitol* 2004; **120**(1): 75–83.
- [27] Itagaki T, Tsutsumi K–i. Triploid form of *Fasciola* in Japan: genetic relationships between *Fasciola hepatica* and *Fasciola gigantica* determined by ITS–2 sequence of nuclear rDNA. *Int J Parasitol* 1998; **28**(5): 777–781.
- [28] Agatsuma T, Arakawa Y, Iwagami M, Honzako Y, Cahyaningsih U, Kang SY, et al. Molecular evidence of natural hybridization between *Fasciola hepatica* and *Fasciola gigantica*. *Parasitol Int* 2000; **49**(3): 231–238.
- [29] Ansari–Lari M, Moazzeni M. A retrospective survey of liver fluke disease in livestock based on abattoir data in Shiraz, south of Iran. *Prev Vet Med* 2006; **73**(1): 93–96.
- [30] Borji H, Parandeh S. The abattoir condemnation of meat because of parasitic infection, and its economic importance: results of a retrospective study in north–eastern Iran. *Ann Trop Med Parasitol* 2010; **104**(8): 641–647.
- [31] Ghazani MH, Valilou MR, Ahmadzadeh AR, Karami AR, Zirak K. The prevalence of sheep liver trematodes in the northwest region of Iran. *Turk J Vet Anim Sci* 2008; **32**(4): 305–307.
- [32] Oryan A, Mansourian M, Moazeni M, Nikahval B, Barband S. Liver distomatosis in cattle, sheep and goats of northeastern Iran. *Global Vet* 2011; **6**(3): 241–246.
- [33] Tavassoli M, Dalirnaghadeh B, Sahandi A. Study on gastrointestinal and external parasites of goats in West Azerbaijan– Iran. *J Vet Res* 2011; **66**(1): 77–79.
- [34] Theodoropoulos G. Risk factors and geospatial modelling for the presence of *Fasciola hepatica* infection in sheep and goat farms in the Greek temperate Mediterranean environment. *Parasitol* 2011; **138**: 926–938.
- [35] Arbabi M, Dalimi A, Ghafarifar F, Foroozandeh MM. Prevalence and intensity of *D. dendriticum* in sheep and goats of Iran. *Res J Parasitol* 2011; **6**(5): 160–167.
- [36] Rojo–Vázquez F, Meana A, Valcárcel F, Martínez–Valladares M. Update on trematode infections in sheep. *Vet Parasitol* 2012; **189**(1): 15–38.
- [37] Cringoli G, Rinaldi L, Veneziano V, Capelli G, Malone JB. A cross–sectional coprological survey of liver flukes in cattle and sheep from an area of the southern Italian Apennines. *Vet Parasitol* 2002; **108**(2): 137–143.
- [38] Otranto D, Traversa D. Dicrocoeliosis of ruminants: a little known fluke disease. *Trends Parasitol* 2003; **19**(1): 12–15.
- [39] Sissay MM, Uggla A, Waller PJ. Prevalence and seasonal incidence of nematode parasites and fluke infections of sheep and goats in eastern Ethiopia. *Trop Anim Health Prod* 2007; **39**(7): 521–531.
- [40] Taylor M. Emerging parasitic diseases of sheep. *Vet Parasitol* 2012; **189**(1): 2–7.
- [41] Ali TS, Zarichehr V, Reza TM, Amroallah B, Hossin T, Amir M, et al. Prevalence of liver flukes infections in slaughtered animals in Kashan, Isfahan province, central Iran. *IIOAB J* 2011; **2**(5): 14–18.
- [42] Khanjari A, Partovi R, Abbaszadeh S, Nemati G, Bahonar A, Misaghi A, et al. A retrospective survey of fasciolosis and dicrocoeliosis in slaughtered animals in Meisam Abattoir, Tehran, Iran (2005–2008). *Vet Res Forum* 2010; **1**(3): 174–178.