



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd



Original article doi: 10.1016/S2222-1808(15)60832-5

©2015 by the Asian Pacific Journal of Tropical Disease. All rights reserved.

Prevalence of intestinal parasitic infections among schoolchildren in Phitsanulok Province, Northern Thailand

Raxsina Polseela, Apichat Vitta*

Department of Microbiology and Parasitology, Faculty of Medical Science, Naresuan University, Phitsanulok, 65000 Thailand

ARTICLE INFO

Article history:

Received 30 Mar 2015

Received in revised form 13 Apr 2015

Accepted 15 May 2015

Available online 22 May 2015

Keywords

Intestinal parasites
Schoolchildren
Phitsanulok Province
Thailand

ABSTRACT

Objective: To determine the prevalence of intestinal parasites in children from six schools located in Bang Rakam and Mueang Districts, Phitsanulok Province, Northern Thailand.**Methods:** The study sites were the following six schools: Wat Krab Phuang, Wat Wang Ped, Wat Tha Ko, Wat Sao Hin, Wat Kung Waree and Wat Sakat Namman. Fecal samples were collected from 352 students in the age range 7-15 years old, and examined for intestinal parasites by formalin-ethyl acetate concentration method. Additionally, Scotch tape technique was used to recover *Enterobius vermicularis* eggs from 576 children.**Result:** It was found that the overall prevalence of intestinal parasites among 352 examined students was 5.4%: *Strongyloides stercoralis* (2.5%), *Entamoeba coli* (*E. coli*) (0.6%), *Giardia lamblia* (0.6%), *Enterobius vermicularis* (0.6%), hookworm (0.3%), hookworm and *E. coli* (0.3%), hookworm and *Trichuris trichiura* (0.3%) and *Strongyloides stercoralis* and *E. coli* (0.3%). The overall prevalence of enterobiasis among 576 schoolchildren by Scotch tape technique was 14.1%. The highest infection rate (19.3%) was observed at Wat Sao Hin School. The selective anthelmintic drugs were administered to all positive cases.**Conclusions:** In this study, low prevalence of parasitic infections was demonstrated among schoolchildren in Phitsanulok Province. However, the control and prevention including the transmission of parasites detected should be explained for more concern and attention and the proper hygienic health education should be implemented.

1. Introduction

Intestinal parasitosis is a major public health problem worldwide, especially among children of developing countries. World Health Organization estimated that in the year 2000 approximately 3.5 billion people all over the world were infected with intestinal parasites and around 450 million children were ill because of these infections. In 1997, it was reported that there were estimated 280 million of hookworm infections worldwide, 478 million of ascariasis and 347 million of trichuriasis cases. *Ascaris lumbricoides* (giant intestinal roundworm), and

Enterobius vermicularis (*E. vermicularis*) (pinworm) were the two common intestinal parasites found in schoolchildren[1,2]. Approximately 320 million schoolchildren worldwide were infected with *Ascaris lumbricoides* and more than 614 million schoolchildren were needed treatment[2,3]. *Enterobius* infection is widespread among children because they are frequently exposed to crowded and inadequate sanitary conditions. Children are particularly vulnerable to infections, causing many nutritional problems, and consequently, resulting in physical and intellectual growth retardation. It is known that intestinal parasitosis can lead to various types of morbidity with the symptoms of bloody stool, chronic diarrhea and abdominal pain. The public health problem of helminthiasis has been neglected in rural areas where there remains a low level of hygiene and poor quality of water supply[4]. Generally, undernourished and weak schoolchildren are also infected with some protozoa with several species inhabited in intestine. Most of them are non-pathogenic

*Corresponding author: Apichat Vitta, Assistant Professor, Department of Microbiology and Parasitology, Faculty of Medical Science, Naresuan University, Phitsanulok, 65000 Thailand.

Tel: 66 055 964653

Fax: 66 055 964770

E-mail: apichatv@nu.ac.th

Foundation Project: Partially supported by Division of Academic Service and Asset Management, Naresuan University.

protozoa such as *Entamoeba coli* (*E. coli*) and *Endolimax nana*, whilst some can cause human symptoms including *Giardia lamblia* (*G. lamblia*) and *Entamoeba histolytica*[5,6].

In Thailand, intestinal parasitosis is the most common infection of people living in rural areas[7,8]. Children are more often infected than adults because they take less care of personal hygiene. Several studies on detection of parasites among schoolchildren in Thailand reported the range of prevalence varied from 4.24% to 75.1%[9-16]. Although the treatments of parasitosis in schoolchildren have been implemented for reducing the parasitic infections, the high prevalence of intestinal parasites in schoolchildren has been widely documented in many areas[9,14-16]. In Phitsanulok Province, few data of parasitic infections among children have been reported[17]. The appropriate control program is necessary to estimate the current status of intestinal parasitic infections. Thus, the objective of this research was to study the prevalence of intestinal parasites among schoolchildren in Phitsanulok Province, Northern Thailand. In addition, Scotch-tape technique was used for detecting *Enterobius* eggs. The results of this study might be useful for the researchers and health authorities for planning and implementing control programs in the studied areas.

2. Materials and methods

The study was performed at six primary schools close to Naresuan University in Phitsanulok Province, Northern Thailand: three schools of Bang Rakam District (Wat Krab Phuang, Wat Wang Ped and Wat Tha Ko) and three schools of Mueang District (Wat Sao Hin, Wat Kung Waree and Wat Sakat Namman) during February 2009 to January 2010. The province is situated in the lower Northern Thailand (Figure 1) and is located 370 km to the north of Bangkok. The majority of people earned their daily lives by agriculture.

An oral description and instruction for handling and collection of fecal samples were given to all children and teachers. The clean containers were distributed to each student on the day before specimen collection. All fecal specimens were transported to the laboratory of Department of Microbiology and Parasitology, Faculty of Medical Science, Naresuan University, and fixed in 10% formalin before further processing. The formalin-ethyl acetate concentration method was used to detect any parasites

presented in fecal samples as described by Garcia[18]. The presence of intestinal parasites was microscopically determined by two parasitologists. In addition, the Scotch tape technique was used for detecting *Enterobius* eggs. The children were explained how to use the Scotch tape slides as described by Beaver *et al.*[19]. All infected children with parasites were treated with anthelmintic drugs and were explained about the transmission of parasites and how to prevent the parasitic diseases.




Figure 1. Map of 6 studied schools in Phitsanulok Province.

3. Results

A total of 352 fecal samples were collected from schoolchildren, 183 boys and 169 girls in the age range from 7-15 years old to check for their intestinal parasites. It was found that the overall prevalence of parasitic infection among schoolchildren was 5.4% (19/352). Among six species of intestinal parasites detected, two species belonged to protozoa (*E. coli* and *G. lamblia*) and other four species were nematodes [*E. vermicularis*, *Trichuris trichiura* (*T. trichiura*), *Strongyloides stercoralis* (*S. stercoralis*), and hookworm]. There were no cestodes and trematodes presented in this study. The infection rate ranged from 0% to 18.7% in different schools, and the highest rate was observed in Wat Wang Ped School. The positive infection rates for boys and girls were 5.5% and 5.3%, respectively. *Strongyloides* infection (2.5%) was the most common (Table 1) whilst *E. coli* and *G. lamblia* were found in 0.6% each. Mixed infections were found between hookworm and *E. coli*, hookworm and *T. trichiura* and *S. stercoralis* and *E. coli* in 0.3%. Hookworm infection and

Table 1

Prevalence of intestinal parasites in schoolchildren of Phitsanulok Province examined by formalin-ethyl acetate concentration technique.

Schools	No. of positive/No. of examined (%)		No. of positive (%)								Total
	Boys	Girls	Ec	Gl	Ev	Hw	Ss	Hw+Ec	Hw+Tt	Ss+Ec	
Wat Krab Phuang	3/45 (6.7)	1/46 (2.2)	2 (2.2)	0	2 (2.2)	0	0	0	0	0	4/91 (4.4)
Wat Kung Waree	0/29 (0.0)	1/23 (4.3)	0	0	0	1 (1.9)	0	0	0	0	1/52 (1.9)
Wat Sakat Namman	1/39 (2.6)	0/37 (0.0)	0	0	0	0	1 (1.3)	0	0	0	1/76 (1.3)
Wat Sao Hin	0/22 (0.0)	0/17 (0.0)	0	0	0	0	0	0	0	0	0/39 (0.0)
Wat Wang Ped	6/34 (17.6)	6/30 (20.0)	0	2 (3.1)	0	0	7 (10.9)	1 (1.6)	1 (1.6)	1 (1.6)	12/64 (18.7)
Wat Tha Ko	0/14 (0.0)	1/16 (6.2)	0	0	0	0	1 (3.3)	0	0	0	1/30 (3.3)
Total	10/183 (5.5)	9/169 (5.3)	2 (0.6)	2 (0.6)	2 (0.6)	1 (0.3)	9 (2.5)	1 (0.3)	1 (0.3)	1 (0.3)	19/ 352 (5.4)

Ec: *E. coli*; Gl: *G. lamblia*; Ev: *E. vermicularis*; Hw: Hookworm; Ss: *S. stercoralis*; Tt: *T. trichiura*.

trichuriasis were individually treated with mebendazole (100 mg) twice daily for three consecutive days.

Additionally, 576 children, 305 boys and 271 girls in the age range of 2-15 years old from four schools, namely, Wat Krab Phuang, Wat Kung Waree, Wat Sakat Namman and Wat Sao Hin were examined by the Scotch tape technique for the presence of *Enterobius* eggs. The overall infection rate was 14.1% (Table 2). The percentage of infected boys and girls was 15.7 and 12.2, respectively. Wat Sao Hin school had the highest infection rate (19.3%). All infected schoolchildren with parasites were individually treated.

Table 2

Prevalence of enterobiasis in schoolchildren of Phitsanulok Province examined by Scotch tape technique.

Schools	No. of positive/No. of examined (%)		Total
	Boys	Girls	
Wat Krab Phuang	14/71 (19.7)	8/70 (11.4)	22/141 (15.6)
Wat Kung Waree	15/73 (20.5)	4/64 (6.2)	19/137 (13.9)
Wat Sakat Namman	13/132 (9.8)	16/109 (14.7)	29/241 (12.0)
Wat Sao Hin	6/29 (20.7)	5/28 (17.8)	11/57 (19.3)
Total	48/305 (15.7)	33/271 (12.2)	81/576 (14.1)

4. Discussion

In the present study, the overall prevalence of intestinal parasites was rather low at 5.4% (19/352) in the studied areas of Phitsanulok Province. Saksirisampant *et al.* (2006) reported that schoolchildren were infected with intestinal parasites at 4.24% (44/1037) in Central Region of Thailand composed of Ang Thong, Ayutthaya and Suphanburi Provinces[12]. In contrast to our study, higher prevalences were reported for Muang Pathum Thani (13.9%)[9], Narathiwat (75%)[11], Ratchaburi (61.1%)[15], Nakhon Prathom (12.6%)[14], and Kanchanaburi (15.6%-23.8%)[16]. The lower prevalence of intestinal parasitosis among schoolchildren in Phitsanulok Province may be due to improvements of general living conditions and the easy accessibility to health care services. High humidity is one factor for parasitic viability in the Southern Thailand where there was high prevalence of soil-transmitted helminths. In addition, migration from Burmese might be one way to make high prevalence of parasites in Ratchaburi and Kanchanaburi. Single parasitic infection was found in 4.55% (16/352) whilst double infection was detected only 0.85% (3/352). Mixed infection of parasites in children was quite common as indicated by Waikagul *et al.*[20]; 33% of schoolchildren from Nan Province were infected with more than one species of parasites.

A non-pathogenic amoeba, *E. coli*, was found in four children, and a pathogenic flagellate protozoa, *G. lamblia*, presented in two children. Generally, these two species were commonly detected in pre-school orphanages and children in primary schools[5,21,22]. However, high infection rates can be also discovered in adults resulting in poor personal and community hygienes. The modes of *Giardia* infection may be due to domestic animals, contaminated food and drinking water.

In the present study, the soil-transmitted helminths found included hookworm, *S. stercoralis* and *T. trichiura* and *S. stercoralis* became the highest detected species with several previous studies at various sites in Thailand and other countries[14,16,22,23]. However, high prevalence of some helminthes such as *T. trichiura* and hookworm was found in Narathiwat Province which was located in the south of Thailand with high moisture and rains throughout the year[11].

The infection rate of enterobiasis among 352 children in six schools examined by formalin-ethyl acetate concentration method was low (0.6%), while the Scotch-tape technique results detected for enterobiasis among 576 children in four schools revealed 14.1%. This technique is suitable for detecting *Enterobius* eggs because the female worms deposit their eggs around anal area.

According to Nithikathkul *et al.* (2001) classification[24], there were three categories of prevalent rates: high (> 27%), intermediate (20%-27%) and low (< 20%); the *Enterobius* prevalence in this study was low consistent with the results observed in other regions of Thailand: 19.7% in kindergarten of Thammasat University[25], and 18.7% in Pichit Province[26]. In contrast, Bunchu and colleagues (2011) reported higher prevalence (25.0%) of enterobiasis in children from five districts in Phitsanulok Province[17], and a high prevalence of this infection (41.6%) was found in children in Mae Suk sub-district in Karen hill tribe villages in Chiang Mai Province[27].

In conclusion, although the overall percentage of intestinal parasitic infection among schoolchildren of Phitsanulok was low, the measures for infection control are recommended for preventing the transmission of parasites. Enterobiasis remains general public health problem among schoolchildren of Phitsanulok Province. An appropriate control of this parasite is needed. Further experiments concerning genotypes and modes of transmission are suggested to perform.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

This study was partially supported by Division of Academic Service and Asset Management, Naresuan University. We would like to thank staff of Department of Microbiology and Parasitology, Faculty of Medical Science, Naresuan University for their technical supports. We also would like to thank Dr. Svetlana Kocherginskaya for editing English grammar in this manuscript.

References

- [1] Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-transmitted helminth infections: ascariasis, trichuriasis,

- and hookworm. *Lancet* 2006; **367**: 1521-32.
- [2] World Health Organization. Eliminating soil-transmitted helminthiasis as a public health problem in children. Geneva: World Health Organization; 2012. [Online] Available from: http://whqlibdoc.who.int/publications/2012/9789241503129_eng.pdf [Accessed on 20 March, 2015]
- [3] World Health Organization. Helminth control in school-age children: a guide for managers of control programmes. Geneva: World Health Organization; 2011. [Online] Available from: http://whqlibdoc.who.int/publications/2011/9789241548267_eng.pdf [Accessed on 20 March, 2015]
- [4] Salam RA, Maredia H, Das JK, Lassi ZS, Bhutta ZA. Community-based interventions for the prevention and control of helminthic neglected tropical diseases. *Infect Dis Poverty* 2014; **3**: 23.
- [5] Lobo ML, Augusto J, Antunes F, Ceita J, Xiao L, Codices V, et al. *Cryptosporidium* spp., *Giardia duodenalis*, *Enterocytozoon bienersi* and other intestinal parasites in young children in Lobata province, Democratic Republic of São Tomé and Príncipe. *PLoS One* 2014; **9**(5): e97708.
- [6] Silva MT, Santana JV, Bragagnoli G, Marinho AM, Malagueño E. Prevalence of *Entamoeba histolytica/Entamoeba dispar* in the city of Campina Grande, in Northeastern Brazil. *Rev Inst Med Trop Sao Paulo* 2014; **56**(5): 451-4.
- [7] Boonjaraspinyo S, Boonmars T, Kaewsamut B, Ekobol N, Laummaunwai P, Aukkanimart R, et al. A cross-sectional study on intestinal parasitic infections in rural communities, Northeast Thailand. *Korean J Parasitol* 2013; **51**(6): 727-34.
- [8] Vitta A, Polseela R, Bunchu N, Waree P, Thongwat D, Tangchaisuriya U, et al. Intestinal helminthiasis in two communities of Phitsanulok Province, Northern Thailand. *J Trop Med Parasitol* 2012; **35**: 1-5.
- [9] Kitvatanachai S, Rhongbutri P. Intestinal parasitic infections in suburban government schools, Lak Hok subdistrict, Muang Pathum Thani, Thailand. *Asian Pac J Trop Med* 2013; **6**(9): 699-702.
- [10] Waikagul J, Jongsuksantigul P, Rattanawitton U, Radomyos P, Kojima S, Takeuchi T. Parasitological monitoring of helminth control program in Northern Thailand. *Southeast Asian J Trop Med Public Health* 2008; **39**: 1008-14.
- [11] Jiraamonninit C, Wongkamchai S, Santabutr W, Loymek S, Monkong N, Nochot H, et al. The prevalence of intestinal parasitic infections among school children with annual anthelmintic treatment in Narathiwat province, Thailand. *J Trop Med Parasitol* 2006; **29**: 45-50.
- [12] Saksirisampant W, Prownebon J, Kulkunthorn M, Yenthakam S, Janpla S, Nuchprayoon S. The prevalence of intestinal parasitic infections among school children in the central region of Thailand. *J Med Assoc Thai* 2006; **89**: 1928-33.
- [13] Wongstitwilairoong B, Srijan A, Serichantalergs O, Fukuda CD, McDaniel P, Bodhidatta L, et al. Intestinal parasitic infections among pre-school children in Sangkhlaburi, Thailand. *Am J Trop Med Hyg* 2007; **76**: 345-50.
- [14] Warunee N, Choomanee L, Sataporn P, Rapeeporn Y, Nuttapong W, Sompong S, et al. Intestinal parasitic infections among school children in Thailand. *Trop Biomed* 2007; **24**: 83-8.
- [15] Maneeboonyang W, Prommongkol S, Wongjindanon N, Treerattanapiboon L, Pasuralertsakul S, Chaimungkun W, et al. Reexamination of parasitic infections in Karen children on the western border of Thailand: two-year follow-up. *J Trop Med Parasitol* 2008; **31**: 77-84.
- [16] Anantaphruti MT, Waikagul J, Maipanich W, Nuamtanong S, Watthanakulpanich D, Pubampen S, et al. School-based health education for the control of soil-transmitted helminthiasis in Kanchanaburi Province, Thailand. *Ann Trop Med Parasitol* 2008; **102**: 521-8.
- [17] Bunchu N, Vitta A, Thongwat D, Lamlerthton S, Pimolsri U, Waree P, et al. *Enterobius vermicularis* infection among children in lower Northern Thailand. *J Trop Med Parasitol* 2011; **34**: 36-40.
- [18] Garcia SL. *Diagnostic medical parasitology*. 5th ed. Washington: ASM press; 2007.
- [19] Beaver PC, Jung RC, Cupp EW. *Clinical parasitology*. 9th ed. Philadelphia: Lea and Febiger; 1984.
- [20] Waikagul J, Krudsood S, Radomyos P, Radomyos B, Chalemrut K, Jonsuksuntigul P, et al. A cross-sectional study of intestinal parasitic infections among school children in Nan Province, Northern Thailand. *Southeast Asian J Trop Med Public Health* 2002; **33**: 218-23.
- [21] Mbae CK, Nokes DJ, Mulinge E, Nyambura J, Waruru A, Kariuk S. Intestinal parasitic infections in children presenting with diarrhoea in outpatient and inpatient settings in an informal settlement of Nairobi, Kenya. *BMC Infect Dis* 2013; **13**: 243.
- [22] Macchioni F, Segundo H, Gabrielli S, Totino V, Gonzales PR, Salazar E, et al. Dramatic decrease in prevalence of soil-transmitted helminths and new insights into intestinal protozoa in children living in the Chaco region, Bolivia. *Am J Trop Med Hyg* 2015; **92**(4): 794-6.
- [23] Karagiannis-Voules DA, Biedermann P, Ekpo UF, Garba A, Langer E, Mathieu E, et al. Spatial and temporal distribution of soil-transmitted helminth infection in sub-Saharan Africa: a systematic review and geostatistical meta-analysis. *Lancet Infect Dis* 2015; **15**(1): 74-84.
- [24] Nithikathkul C, Changsap B, Wannapinyosheep S, Poister C, Boontan P. The prevalence of enterobiasis in children attending mobile health clinic of Huachiew Chalermprakiet University. *Southeast Asian J Trop Med Public Health* 2001; **32**(Suppl 2): 138-42.
- [25] Pethleart A, Saichua P, Rhongbutri P, Leelawongtawon R, Aree K, Tiengtip R, et al. Prevalence and risk factors for pinworm infection in the kindergarten of Thammasat University, Thailand. *Southeast Asian J Trop Med Public Health* 2010; **41**: 306-10.
- [26] Nateeworanart S, Vitta A, Lee UP. Egg positive rate of *Enterobius vermicularis* in children in a rural area of Phichit province, Thailand. *Southeast Asian J Trop Med Public Health* 2007; **38**(Suppl 1): 40-2.
- [27] Chaisalee T, Tukaew A, Suwansakri J. Very high prevalence of enterobiasis among the hilltribal children in rural district "Mae Suk" Thailand. *Med Gen Med* 2004; **6**: 5.